

On the properties of circulant matrices involving Generalized Tribonacci and Generalized Tribonacci-Lucas numbers

Nazmiye Yilmaz*

Department of Mathematics, Science Faculty,
Selcuk University, Campus, Konya, Turkey

Yasin Yazlik

Department of Mathematics, Faculty of Science and Letters,
Nevsehir Hacı Bektas Veli University, Nevsehir, Turkey

Necati Taskara

Department of Mathematics, Science Faculty,
Selcuk University, Campus, Konya, Turkey

July 18, 2014

Abstract

In this paper, firstly, we define the Generalized Tribonacci-Lucas numbers. In addition, by also defining circulant matrices $C_n(G)$ and $C_n(S)$ whose entries are Generalized Tribonacci and Generalized Tribonacci-Lucas numbers, we compute spectral norms and determinants of these matrices.

Keywords: Generalized Tribonacci numbers, Generalized Tribonacci-Lucas numbers, circulant matrix, norm, determinant.

AMS Classification: 11B39, 15A18, 15A60.

1 Introduction

Shannon and Horadam [5] defined Generalized Tribonacci sequence $\{G_n\}_{n \in \mathbb{N}}$ as

$$G_{n+3} = PG_{n+2} + QG_{n+1} + RG_n, \quad G_0 = 0, \quad G_1 = 0, \quad G_2 = 1, \quad (1)$$

where P , Q , and R are arbitrary integers.

If the characteristic equation

$$x^3 - Px^2 - Qx - R = 0$$

*e mail: nzyilmaz@selcuk.edu.tr, yyazlik@nevsehir.edu.tr, ntaskara@selcuk.edu.tr

has three distinct real roots, suppose that they are given by α, β, γ . According to the general theory of recurrence relations, G_n can be represented by

$$G_n = \frac{\alpha^n}{(\alpha - \beta)(\alpha - \gamma)} + \frac{\beta^n}{(\beta - \alpha)(\beta - \gamma)} + \frac{\gamma^n}{(\gamma - \alpha)(\gamma - \beta)}. \quad (2)$$

The following relations hold:

$$\alpha + \beta + \gamma = P, \quad \alpha\beta\gamma = R, \quad \alpha\beta + \beta\gamma + \alpha\gamma = -Q, \quad (3)$$

where α, β and γ are roots of equation (1).

Recently, the norms of some special matrices were studied. For instance, Solak [7], defined the $n \times n$ circulant matrices with Fibonacci and Lucas numbers entries. Additionally, he investigated the upper and lower bounds of spectral norms of the circulant matrices. Shen and Cen, in [6], have found upper and lower bounds for the spectral norms of r -circulant matrices and obtained some bounds for the spectral norms of Kronecker and Hadamard products of these matrices. In [4], the norms, eigenvalues and determinants of some matrices related with special numbers were studied. In [3], Ipek obtained the spectral norms of circulant matrices with Fibonacci and Lucas numbers entries. In [8], the authors computed the spectral norm, eigenvalues and determinant of circulant matrix involving the Generalized k -Horadam numbers.

Here, we define the Generalized Tribonacci-Lucas numbers and present eigenvalues, spectral norms and determinants of circulant matrices with Generalized Tribonacci and Generalized Tribonacci-Lucas numbers.

2 Circulant Matrix with Generalized Tribonacci numbers

From [1], we first remind that the circulant matrix $C = [c_{ij}] \in M_{n,n}(\mathbb{C})$ is defined by

$$c_{ij} = \begin{cases} c_{j-i}, & j \geq i \\ c_{n+j-i}, & j < i \end{cases}.$$

From [2], we further remind that, for a matrix $X = [x_{i,j}] \in M_{m,n}(\mathbb{C})$, the spectral norm of X is given by

$$\|X\|_2 = \sqrt{\max_{1 \leq i \leq n} \lambda_i(X^*X)}$$

where X^* is the conjugate transpose of matrix X .

In this section, we define eigenvalues, the spectral norm and determinant of the circulant matrix with the Generalized Tribonacci numbers entries. In order to do that, we can also define *the circulant matrix* as follows.

Definition 1 A $n \times n$ circulant matrix with Generalized Tribonacci numbers entries is defined by

$$C_n(G) = \begin{bmatrix} G_1 & G_2 & G_3 & \cdots & G_n \\ G_n & G_1 & G_2 & \cdots & G_{n-1} \\ G_{n-1} & G_n & G_1 & \cdots & G_{n-2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ G_2 & G_3 & G_4 & \cdots & G_1 \end{bmatrix}. \quad (4)$$

Lemma 2 [1] Let $A = \text{circ}(a_0, a_1, \dots, a_{n-1})$ be a $n \times n$ circulant matrix. Then we have

$$\lambda_j(A) = \sum_{k=0}^{n-1} a_k w^{-jk},$$

where $w = e^{\frac{2\pi i}{n}}$, $i = \sqrt{-1}$, $j = 0, 1, \dots, n-1$.

Lemma 3 [2] Let A be an $n \times n$ matrix with eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$. Then, A is a normal matrix if and only if the eigenvalues of AA^* are $|\lambda_1|^2, |\lambda_2|^2, \dots, |\lambda_n|^2$, where A^* is the conjugate transpose of the matrix A .

The following result gives us the eigenvalues of the matrix in (4).

Theorem 4 The eigenvalues of $C_n(G)$ are

$$\lambda_j(C_n(G)) = \frac{G_{n+1} + (G_{n+2} - PG_{n+1} - 1)w^{-j} + RG_n w^{-2j}}{Pw^{-j} + Qw^{-2j} + Rw^{-3j} - 1},$$

where G_n is the n th Generalized Tribonacci number and P , Q , and R are arbitrary integers.

Proof. By considering $X = (\alpha - \beta)(\alpha - \gamma)$, $Y = (\beta - \alpha)(\beta - \gamma)$, $Z = (\gamma - \alpha)(\gamma - \beta)$, from Lemma 2 and (2), we have

$$\begin{aligned} \lambda_j(C_n(G)) &= \sum_{k=0}^{n-1} G_{k+1} w^{-jk} = \sum_{k=0}^{n-1} \left(\frac{\alpha^{k+1}}{X} + \frac{\beta^{k+1}}{Y} + \frac{\gamma^{k+1}}{Z} \right) w^{-jk} \\ &= \frac{\alpha}{X} \left(\frac{(\alpha w^{-j})^n - 1}{\alpha w^{-j} - 1} \right) + \frac{\beta}{Y} \left(\frac{(\beta w^{-j})^n - 1}{\beta w^{-j} - 1} \right) + \frac{\gamma}{Z} \left(\frac{(\gamma w^{-j})^n - 1}{\gamma w^{-j} - 1} \right) \\ &= \frac{\alpha^{n+1} - \alpha}{X(\alpha w^{-j} - 1)} + \frac{\beta^{n+1} - \beta}{Y(\beta w^{-j} - 1)} + \frac{\gamma^{n+1} - \gamma}{Z(\gamma w^{-j} - 1)}. \end{aligned}$$

By using (2) and (3), we obtain

$$\lambda_j(C_n(G)) = \frac{G_{n+1} + (G_{n+2} - PG_{n+1} - 1)w^{-j} + RG_n w^{-2j}}{Pw^{-j} + Qw^{-2j} + Rw^{-3j} - 1},$$

as required. ■

The following theorem gives us the spectral norm of $C_n(G)$.

Theorem 5 For matrix in (4), we have

$$\|C_n(G)\|_2 = \frac{G_{n+2} + (1-P)G_{n+1} + RG_n - 1}{P + Q + R - 1},$$

where P , Q , and R are arbitrary integers and $P + Q + R - 1 \neq 0$.

Proof. We know that matrix $C_n(G)$ in (4) is a normal matrix since $C_n(G)^*C_n(G) = C_n(G)C_n(G)^*$. Now, from Lemma 3, we can write

$$\|C_n(G)\|_2 = \sqrt{\left(\max_{0 \leq j \leq n-1} |\lambda_j(C_n(G))|^2\right)}.$$

In this last equality, if we take $j = 0$, then λ_0 becomes the maximum eigenvalue. Thus, $\|C_n(G)\|_2 = |\lambda_0(C_n(G))|$. In addition, Theorem 4, we clearly obtain

$$\|C_n(G)\|_2 = \frac{G_{n+2} + (1-P)G_{n+1} + RG_n - 1}{P + Q + R - 1}.$$

Hence, proof is completed. ■

In particular cases:

- Taking $P = Q = R = 1$, then we get $\|C_n(T)\|_2 = \frac{T_{n+1} + T_{n-1} - 1}{2}$, where T_n is n th the classical Tribonacci number.
- Taking $P = 0$, $Q = R = 1$, then we get $\|C_n(A)\|_2 = A_{n+1} - 1$, where A_n is n th the Padovan number.
- Taking $P = Q = 1$, $R = 0$, then we get $\|C_n(F)\|_2 = F_{n+1} - 1$ in [3], where F_n is n th the classical Fibonacci number.
- Taking $P = k$, $Q = 1$, $R = 0$, then we get $\|C_n(F_k)\|_2 = \frac{F_{k,n} + F_{k,n-1} - 1}{k}$, where $F_{k,n}$ is n th the k -Fibonacci number, that is $F_{k,n} = kF_{k,n-1} + F_{k,n-2}$ with initial conditions $F_{k,0} = 0$ and $F_{k,1} = 1$.
- Taking $P = 2$, $Q = 1$, $R = 0$, then we get $\|C_n(B)\|_2 = \frac{B_{n+1} - B_{n-1}}{2}$, where B_n is n th the classical Pell number.
- Taking $P = 1$, $Q = 2$, $R = 0$, then we get $\|C_n(J)\|_2 = \frac{J_{n+1} - 1}{2}$, where J_n is n th the Jacobsthal number.

Theorem 6 The determinant of the matrix $C_n(G) = \text{circ}(G_1, G_2, \dots, G_n)$ is written by

$$\det(C_n(G)) = \frac{(G_{n+1})^n (1 - K^n - L^n + K^n L^n)}{(-1)^n (1 - S_{n+1} - R^n + M)},$$

where

$$\begin{aligned} K &= \frac{PG_{n+1} - G_{n+2} + 1 - \sqrt{(PG_{n+1} - G_{n+2} + 1)^2 - 4RG_nG_{n+1}}}{2G_{n+1}}, \\ L &= \frac{PG_{n+1} - G_{n+2} + 1 + \sqrt{(PG_{n+1} - G_{n+2} + 1)^2 - 4RG_nG_{n+1}}}{2G_{n+1}}, \\ M &= \alpha^n \beta^n + \beta^n \gamma^n + \alpha^n \gamma^n. \end{aligned}$$

Proof. From Theorem 4, we have

$$\begin{aligned} \det(C_n(G)) &= \prod_{j=0}^{n-1} \lambda_j(C_n(G)) \\ &= \prod_{j=0}^{n-1} \frac{G_{n+1} - (PG_{n+1} - G_{n+2} + 1)w^{-j} + RG_nw^{-2j}}{(\alpha w^{-j} - 1)(\beta w^{-j} - 1)(\gamma w^{-j} - 1)}. \end{aligned}$$

By considering identities $\prod_{k=0}^{n-1} (x - yw^{-k}) = x^n - y^n$ and

$$\prod_{k=0}^{n-1} (x - yw^{-k} + zw^{-2k}) = x^n \left(1 - \left(\frac{y - \sqrt{y^2 - 4xz}}{2x} \right)^n - \left(\frac{y + \sqrt{y^2 - 4xz}}{2x} \right)^n + \left(\frac{y - \sqrt{y^2 - 4xz}}{2x} \right)^n \left(\frac{y + \sqrt{y^2 - 4xz}}{2x} \right)^n \right),$$

we can write

$$\det(C_n(G)) = \frac{(G_{n+1})^n (1 - K^n - L^n + K^n L^n)}{(-1)^n (1 - S_{n+1} - R^n + M)}.$$

■

3 Circulant Matrix with Generalized Tribonacci-Lucas numbers

In this section, we firstly define Generalized Tribonacci-Lucas sequence $\{S_n\}_{n \in \mathbb{N}}$. Then we compute the eigenvalues, norm and determinant of circulant matrix with the Generalized Tribonacci-Lucas sequence $\{S_n\}_{n \in \mathbb{N}}$.

Definition 7 For $R \neq 0$, the Generalized Tribonacci-Lucas sequence $\{S_n\}_{n \in \mathbb{N}}$ is defined by

$$S_{n+3} = PS_{n+2} + QS_{n+1} + RS_n, \quad S_0 = -\frac{Q}{R}, \quad S_1 = 3, \quad S_2 = P. \quad (5)$$

Binet formula enables us to state the Generalized Tribonacci-Lucas numbers. It can be clearly obtained from the three distinct real roots α , β and γ of characteristic equation of (5) as

$$S_n = \alpha^{n-1} + \beta^{n-1} + \gamma^{n-1}. \quad (6)$$

Note that, α , β and γ are the same in (3).

Definition 8 The $n \times n$ circulant matrix with the Generalized Tribonacci numbers entries is defined by

$$C_n(S) = \begin{bmatrix} S_1 & S_2 & S_3 & \cdots & S_n \\ S_n & S_1 & S_2 & \cdots & S_{n-1} \\ S_{n-1} & S_n & S_1 & \cdots & S_{n-2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ S_2 & S_3 & S_4 & \cdots & S_1 \end{bmatrix}. \quad (7)$$

The following result gives us the eigenvalues of the matrix in (7).

Theorem 9 For $w = e^{\frac{2\pi i}{n}}$, $i = \sqrt{-1}$ and $j = 0, 1, \dots, n-1$, the eigenvalues of $C_n(S)$ are

$$\lambda_j(C_n(S)) = \frac{S_{n+1} - 3 + (S_{n+2} - PS_{n+1} + 2P)w^{-j} + (RS_n + Q)w^{-2j}}{Pw^{-j} + Qw^{-2j} + Rw^{-3j} - 1},$$

where S_n is the n th Generalized Tribonacci-Lucas number and P , Q , and R are arbitrary integers.

Proof. By considering Lemma 2 and (6), we have

$$\begin{aligned} \lambda_j(C_n(S)) &= \sum_{k=0}^{n-1} S_{k+1} w^{-jk} = \sum_{k=0}^{n-1} (\alpha^k + \beta^k + \gamma^k) w^{-jk} \\ &= \frac{\alpha^n - 1}{\alpha w^{-j} - 1} + \frac{\beta^n - 1}{\beta w^{-j} - 1} + \frac{\gamma^n - 1}{\gamma w^{-j} - 1}. \end{aligned}$$

By using (3), we obtain

$$\lambda_j(C_n(S)) = \frac{S_{n+1} - 3 + (S_{n+2} - PS_{n+1} + 2P)w^{-j} + (RS_n + Q)w^{-2j}}{Pw^{-j} + Qw^{-2j} + Rw^{-3j} - 1}.$$

■

Theorem 10 For the matrix in (7), the following equalities hold:

$$\text{i) } \|C_n(S)\|_2 = \frac{S_{n+2} + (1-P)S_{n+1} + RS_n + 2P + Q - 3}{P + Q + R - 1},$$

where P , Q , and R are arbitrary integers and $P + Q + R - 1 \neq 0$.

$$\text{ii) } \det(C_n(S)) = \frac{(S_{n+1}-3)^n [1-C^n-D^n+C^n D^n]}{(-1)^n [1-S_{n+1}-R^n+E]},$$

where

$$\begin{aligned} C &= \frac{S_{n+2} - PS_{n+1} + 2P - \sqrt{(S_{n+2} - PS_{n+1} + 2P)^2 - 4(S_{n+1} - 3)(RS_n + Q)}}{2(S_{n+1} - 3)}, \\ D &= \frac{S_{n+2} - PS_{n+1} + 2P + \sqrt{(S_{n+2} - PS_{n+1} + 2P)^2 - 4(S_{n+1} - 3)(RS_n + Q)}}{2(S_{n+1} - 3)}, \\ E &= \alpha^n \beta^n + \beta^n \gamma^n + \alpha^n \gamma^n. \end{aligned}$$

Proof. Proof of this theorem can be seen by using Theorem 9. ■

In particular cases:

- Taking $P = Q = R = 1$, then we get $\|C_n(Y)\|_2 = \frac{Y_{n+1}+Y_{n-1}}{2}$, where Y_n is n th the classical Tribonacci-Lucas number.
- Taking $P = 0$, $Q = R = 1$, then we get $\|C_n(Z)\|_2 = Z_{n+4} - 2$, where Z_n is n th the Perrin number.

References

- [1] P. J. Davis, Circulant Matrices (John Wiley&Sons, New York, 1979).
- [2] R. A. Horn, C. R. Johnson, Matrix Analysis (Cambridge University Press, 1985).
- [3] A. Ipek, On the spectral norms of circulant matrices with classical Fibonacci and Lucas numbers entries, *Applied Mathematics and Computation*, **217**(12), (2011), 6011-6012.
- [4] E.G. Kocer, Circulant, negacyclic and semicirculant matrices with the modified Pell, Jacobsthal and Jacobsthal-Lucas numbers, *Hacettepe J. Math., and Statistics*, **36**(2), (2007), 133-142.
- [5] A.G. Shanon, A.F. Horadam, Some properties of third-order recurrence relations, *Fibonacci Quarterly*, **10**(2), (1972), 135-145.
- [6] S. Shen, J. Cen, On the bounds for the norms of r -circulant matrices with the Fibonacci and Lucas numbers, *Applied Mathematics and Computation*, **216**, (2010), 2891-2897.
- [7] S. Solak, On the norms of circulant matrices with the Fibonacci and Lucas numbers, *Applied Mathematics and Computation*, **160**, (2005), 125-132.
- [8] Y. Yazlik, N. Taskara, Spectral norm, Eigenvalues and Determinant of Circulant Matrix involving the Generalized k -Horadam numbers, *Ars Combinatoria*, **104**, (2012), 505-512.